



Site Name: Lough Corrib SAC

Site Code: 000297

Lough Corrib is situated to the north of Galway city and is the second largest lake in Ireland, with an area of approximately 18,240 ha (the entire site is 20,556 ha). The lake can be divided into two parts: a relatively shallow basin, underlain by Carboniferous limestone, in the south, and a larger, deeper basin, underlain by more acidic granite, schists, shales and sandstones to the north. The surrounding lands to the south and east are mostly pastoral farmland, while bog and heath predominate to the west and north. A number of rivers are included within the cSAC as they are important for Atlantic Salmon. These rivers include the Clare, Grange, Abbert, Sinking, Dalgan and Black to the east, as well as the Cong, Bealanabrack, Failmore, Cornamona, Drimneen and Owenriff to the west. In addition to the rivers and lake basin, adjoining areas of conservation interest, including raised bog, woodland, grassland and limestone pavement, have been incorporated into the site.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

- [3110] Oligotrophic Waters containing very few minerals
- [3140] Hard Water Lakes
- [3260] Floating River Vegetation
- [6210] Orchid-rich Calcareous Grassland*
- [6410] *Molinia* Meadows
- [7110] Raised Bog (Active)*
- [7120] Degraded Raised Bog
- [7150] Rhynchosporion Vegetation
- [7210] *Cladium* Fens*
- [7220] Petrifying Springs*
- [7230] Alkaline Fens
- [8240] Limestone Pavement*
- [91A0] Old Oak Woodlands
- [91D0] Bog Woodland*

- [1029] Freshwater Pearl Mussel (*Margaritifera margaritifera*)
- [1092] White-clawed Crayfish (*Austropotamobius pallipes*)
- [1095] Sea Lamprey (*Petromyzon marinus*)
- [1096] Brook Lamprey (*Lampetra planeri*)

- [1106] Atlantic Salmon (*Salmo salar*)
- [1303] Lesser Horseshoe Bat (*Rhinolophus hipposideros*)
- [1355] Otter (*Lutra lutra*)
- [1393] Slender Green Feather-moss (*Drepanocladus vernicosus*)
- [1833] Slender Naiad (*Najas flexilis*)

The shallow, lime-rich waters of the southern basin of Lough Corrib support one of the most extensive beds of stoneworts (Charophytes) in Ireland, with species such as *Chara aspera*, *C. hispida*, *C. delicatula*, *C. contraria* and *C. desmacantha* mixed with submerged pondweeds (*Potamogeton perfoliatus*, *P. gramineus* and *P. lucens*), Shoreweed (*Littorella uniflora*) and Water Lobelia (*Lobelia dortmanna*). These *Chara* beds are an important source of food for waterfowl. In contrast, the northern basin contains more oligotrophic and acidic waters, without *Chara* species, but with Shoreweed, Water Lobelia, Pipewort (*Eriocaulon aquaticum*), Quillwort (*Isoetes lacustris*), Alternate Water-milfoil (*Myriophyllum alternifolium*) and Slender Naiad (*Najas flexilis*). The last-named is listed under the Flora (Protection) Order, 1999, and is an Annex II species under the E.U. Habitats Directive.

Large areas of reedswamp vegetation, dominated by varying mixtures of Common Reed (*Phragmites australis*) and Common Club-rush (*Scirpus lacustris*), occur around the margins of the lake. Reedswamp usually grades into species-rich marsh vegetation characterised by Slender Sedge (*Carex lasiocarpa*), Water Mint (*Mentha aquatica*), Water Horsetail (*Equisetum fluviatile*) and Bogbean (*Menyanthes trifoliata*). Of particular note are the extensive beds of Great Fen-sedge (*Cladium mariscus*) that have developed over the marly peat deposits in sheltered bays, particularly in the south-east corner of the lake. Alkaline fen vegetation is more widespread around the lake margins and includes, amongst the typically diverse range of plants, the Slender Cottongrass (*Eriophorum gracile*), a species protected under the Flora (Protection) Order, 1999. Wet meadows dominated by Purple Moor-grass (*Molinia caerulea*) occur in seasonally flooded areas close to the lake shore. These support species such as Sharp-flowered Rush (*Juncus acutiflorus*), Jointed Rush (*J. articulatus*), Carnation Sedge (*Carex panicea*), Devil's-bit Scabious (*Succisa pratensis*), Creeping Bent (*Agrostis stolonifera*) and Tormentil (*Potentilla erecta*), amongst others.

This large site contains four discrete raised bog areas and is selected for active raised bog, degraded raised bog, Rhynchosporion and bog woodland. Active raised bog comprises areas of high bog that are wet and actively peat-forming, where the percentage cover of bog mosses (*Sphagnum* spp.) is high, and where some or all of the following features occur: hummocks, pools, wet flats, *Sphagnum* lawns, flushes and soaks. Degraded raised bog corresponds to those areas of high bog whose hydrology has been adversely affected by peat cutting, drainage and other land use activities, but which are capable of regeneration. The Rhynchosporion habitat occurs in wet depressions, pool edges and erosion channels where the vegetation includes White Beak-sedge (*Rhynchospora alba*) and/or Brown Beak-sedge (*R. fusca*), and at least some of the following associated species, Bog Asphodel (*Narthecium ossifragum*), sundews (*Drosera* spp.), Deergrass (*Scirpus cespitosus*) and Carnation Sedge.

At Addergoole, on the eastern shores of Lough Corrib, there is an important area of western raised bog. This bog area is one of the most westerly, relatively intact raised bogs in the country. There are also other substantial areas of raised bog along various tributaries of the Corrib in east Co. Galway, namely Slieve Bog, Lough Tee Bog and Killaclogher bog. The active parts of these bogs mostly correspond to the wettest areas, where there are well-developed surface features with hummocks, lawns and pools. It is in such areas that Rhynchosporion vegetation is best represented. The dominant species is the aquatic bog moss *Sphagnum cuspidatum*, which is usually accompanied by Bogbean, White Beak-sedge, Bog Asphodel, Common Cottongrass (*Eriophorum angustifolium*), Bog Sedge (*Carex limosa*) and Great Sundew (*Drosera anglica*). Brown Beak-sedge, a locally rare plant of wet bog pools, has been recorded from a number of the bog areas within the site. At Addergoole a substantial bog lake or soak occurs and this is infilling with large rafts of Rhynchosporion vegetation at present. This area is associated with an important area of wet bog woodland dominated by Downy Birch (*Betula pubescens*).

The largest part of the uncut high bog comprises degraded raised bog. Degraded bog is dominated by a raised bog flora which tends to be rather species-poor because of disturbance and/or drying-out. The most conspicuous vascular plant species are usually Carnation Sedge, Heather (*Calluna vulgaris*), Cottongrasses, Cross-leaved Heath (*Erica tetralix*), Bog Asphodel and Deergrass. Bog-rosemary (*Andromeda polifolia*) and Cranberry (*Vaccinium oxycoccos*), two species indicative of raised bog habitat, are frequent on both degraded and active areas of raised bog. *Sphagnum* cover is generally low within degraded areas due to a combination of drying-out and frequent burning.

Limestone pavement occurs along much of the shoreline in the lower Corrib basin, and supports a rich and diverse flora, including Herb-Robert (*Geranium robertianum*), Bloody Crane's-bill (*G. sanguineum*), Carline Thistle (*Carlina vulgaris*), Spring Gentian (*Gentiana verna*), Wild Thyme (*Thymus praecox*), Rustyback (*Ceterach officinarum*), Wood Sage (*Teucrium scorodonia*), Slender St. John's-wort (*Hypericum pulchrum*), Quaking-grass (*Briza media*) and Blue Moor-grass (*Sesleria albicans*). Areas of Hazel (*Corylus avellana*) scrub occur in association with exposed limestone pavement and these include species such as Hawthorn (*Crataegus monogyna*), Buckthorn (*Rhamnus catharticus*), Spindle (*Euonymus europaeus*), with occasional Juniper (*Juniperus communis*). Three Red Data Book species are also found in association with limestone scrub - Alder Buckthorn (*Frangula alnus*), Shrubby Cinquefoil (*Potentilla fruticosa*) and Wood Bitter-vetch (*Vicia orobus*), the latter is also protected under the Flora (Protection) Order, 1999.

Open areas of orchid-rich calcareous grassland are also found in association with the limestone exposures. These can support a typically rich vegetation, including many orchids such as Pyramidal Orchid (*Anacamptis pyramidalis*), Common Spotted-orchid (*Dactylorhiza fuchsii*), Early-purple Orchid (*Orchis mascula*), Frog Orchid (*Coeloglossum viride*), Fragrant Orchid (*Gymnadenia conopsea*), Marsh Helleborine (*Epipactis palustris*), Greater Butterfly-orchid (*Platanthera chlorantha*) and Irish Lady's-tresses

(*Spiranthes romanzoffiana*). The latter is protected under the Flora (Protection) Order, 1999.

The Hill of Doon, located in the north-western corner of the lake, is a fine example of a Sessile Oak (*Quercus petraea*) woodland. The understorey is dominated by Sessile Oak, Holly (*Ilex aquifolium*) and occasional Juniper. There are occasional Yew (*Taxus baccata*) and Ash (*Fraxinus excelsior*), and a well developed ground layer dominated by Bilberry (*Vaccinium myrtillus*), Hard Fern (*Blechnum spicant*) and Wood Rush (*Luzula sylvatica*). Woodland also occurs on some of the islands in the lake.

A number of the rivers in the site support submerged and floating vegetation of the Ranunculion fluitantis and Callitriche-Batrachion, including mosses. For example, in the River Corrib species such as Shining Pondweed (*Potamogeton lucens*), Perfoliate Pondweed (*Potamogeton perfoliatus*), Small Pondweed (*P. berchtoldii*), Yellow Water-lily (*Nuphar lutea*), White Water-lily (*Nymphaea alba*) and stoneworts (*Chara* spp.) occur.

The rare and Annex II-listed Slender Green Feather-moss (*Drepanocladus vernicosus*) is found at the fen at Gortachalla, north-east of Moycullen. Here it is widespread around the margins, and this constitutes a large and significant population in the national context. A very large population of another rare moss, *Pseudocalliergon trifarium*, is also found in this area.

The lake is rated as an internationally important site for waterfowl. Counts from 1984 to 1987 revealed a mean annual peak total of 19,994 birds. In the past a maximum peak of 38,281 birds was recorded. The lake supports internationally important numbers of Pochard (average peak 8,600) and nationally important numbers of the following species: Coot (average peak 6,756), Mute Swan (average peak 176), Tufted Duck (average peak 1,317), Cormorant (average peak 110) and Greenland White-fronted Goose (average peak 83). The latter species is listed on Annex I of the E.U. Birds Directive. The Coot population is the largest in the country and populations of Tufted Duck and Pochard are second only to Lough Neagh. Breeding pairs of Common Scoter on the lake number 30-41 (1995 data), as well as breeding populations of Arctic Tern and Common Tern. Other bird species of note recorded from or close to the lake recently include Hen Harrier, Whooper Swan, Golden Plover and Kingfisher. All of these species are listed on Annex I of the E.U. Birds Directive.

Otter and Irish Hare have been recorded regularly within this site. Both of these species are listed in the Red Data Book and are legally protected by the Wildlife Act, 1976. Otter is also listed on Annex II of the E.U. Habitats Directive. Lough Corrib is considered one of the best sites in the country for Otter, due to the sheer size of the lake and associated rivers and streams, and also the generally high quality of the habitats. Atlantic Salmon (*Salmo salar*) use the lake and rivers as spawning grounds. Although this species is still fished commercially in Ireland, it is considered to be endangered or locally threatened elsewhere in Europe and is listed on Annex II of the E.U. Habitats Directive. Lough Corrib is also a well known fishing lake with a

very good Trout (*Salmo trutta*) fishery. The lake has a population of Sea Lamprey (*Petromyzon marinus*), a scarce, though probably under-recorded species listed on Annex II of the E.U. Habitats Directive. Brook Lamprey (*Lampetra planeri*), also listed on Annex II, are also known from a number of areas within the site.

A population of Freshwater Pearl Mussel (*Margaritifera margaritifera*), a species listed on Annex II of the E.U. Habitats Directive, occurs within the site. White-clawed Crayfish (*Austropotamobius pallipes*), also listed on Annex II, is well distributed throughout Lough Corrib and its in-flowing rivers over limestone. A summer roost of Lesser Horseshoe Bat, another Annex II species, occurs within the site - approximately 100 animals were recorded here in 1999.

The main threats to the quality of this site are from water polluting activities resulting from intensification of agricultural activities on the eastern side of the lake, uncontrolled discharge of sewage which is causing localised eutrophication of the lake, and housing and boating development, which is causing the loss of native lakeshore vegetation. The raised bog habitats are susceptible to further degradation and drying out due to drainage and peat cutting and, on occasions, burning. Peat cutting threatens Addergoole Bog and already a substantial area of it has been cut away. Fishing and shooting occur in and around the lake. Introduction of exotic crayfish species or the crayfish fungal plague (*Aphanomyces astaci*) could have a serious impact on the native crayfish population. The bat roost is susceptible to disturbance or development.

Despite these ongoing issues, however, Lough Corrib is one the best examples of a large lacustrine catchment system in Ireland, with a range of habitats and species still well represented. These include 14 habitats which are listed on Annex I of the E.U. Habitats Directive, six of which are priority habitats, and nine species which are listed on Annex II. The lake is also internationally important for birds and is designated as a Special Protection Area.